


PART 2

Regulation of IL-6 Signaling

The correct response of the target cell to cytokines is crucial

Serge Haan, Habilitation 2005





Outline: Interleukin-6 signal transduction and its regulation

Part 1: Molecular mechanisms of IL-6 signal transduction

Part 2: Regulation of IL-6 signal transduction




- **Half-lives of the signaling components**
- Polar expression of IL-6 receptor- α (gp80)
- IL-6 receptor- α shedding
- Internalization of the ligand/IL-6 receptor complex (desensitisation)
- Phosphotyrosine-phosphatase SHP2 deactivates receptors, JAKs and STAT3
- SOCS proteins (suppressors of cytokine signaling) are feedback inhibitors
- Protein inhibitors of activated STATs (PIAS)
- Cross-talks between JAK/STAT signaling and proinflammatory cytokines (IL-1, TNF α)
 - STAT3 and NF- κ B compete for an overlapping response element
 - IL-1 stabilizes SOCS3-mRNA
 - IL-1 accelerates the internalization of the signal transducer gp130

Structural organization of various IL-6-type cytokine signaling components



CBM	Cytokine Binding Module
ESS	Extended SH2 Subdomain
FERM	4.1/Ezrin/Radixin/Moesin
FNIII	Fibronectin type III
KIR	Kinase Inhibitory Region
PTB	Phosphotyrosine Binding
SH2	Src Homology 2
TM	Transmembrane

Half-lives of proteins involved in IL-6-type cytokine signaling



COS-7 cells were pulse-labelled with [35 S]methionine/cysteine and chased for the times indicated (A, 0.25±12 h; B and C, 0.5±72 h).

Calculated half-lives of IL-6 signaling components

Protein	Half-life [h]	Protein	Half-life [h]
gp80 (IL-6R α)	2.5	STAT3 α	8.5
gp130	2.5	STAT3 β	4.5
Jak1	3.2	STAT3 Δ 715	8.5
Jak2	1.9		
Tyk2	2.0		
SOCS1	1.5	STAT1	16.0
SOCS2	1.0	SHP2	18.0-20.0
SOCS3	1.6		

STAT3 β is a splice variant of STAT3 α lacking the 55 C-terminal amino acids which are substituted by seven STAT3 β -specific amino acids.

STAT3 Δ 715 = STAT3 deletion mutant

Gerhartz et al. Eur. J. Biochem. 223, 265-274 (1994) [48 citations](#) (07/2022)

Siewert et al. Eur. J. Biochem. 265, 251-257 (1999) [93 citations](#) (07/2022)


Outline: Interleukin-6 signal transduction and its regulation

Part 1: Molecular mechanisms of IL-6 signal transduction


Part 2: Regulation of IL-6 signal transduction

- Half-lives of the signaling components
- **Polar expression of IL-6 receptor- α (gp80)**
- IL-6 receptor- α shedding
- Internalization of the ligand/IL-6 receptor complex (desensitisation)
- Phosphotyrosine-phosphatase SHP2 deactivates receptors, JAKs and STAT3
- SOCS proteins (suppressors of cytokine signaling) are feedback inhibitors
- Protein inhibitors of activated STATs (PIAS)
- Cross-talks between JAK/STAT signaling and proinflammatory cytokines (IL-1, TNF α)
 - STAT3 and NF- κ B compete for an overlapping response element
 - IL-1 stabilizes SOCS3-mRNA
 - IL-1 accelerates the internalization of the signal transducer gp130


IL-6 receptor α -chain:
What is the function of the cytoplasmic part?



IL-6 receptor α chain:
What is the function of the cytoplasmic part?



IL-6 receptor α chain:
What is the function of the cytoplasmic part?



The cytoplasmic domain of the interleukin-6 receptor gp80 mediates its basolateral sorting in polarized Madin-Darby canine kidney cells

25 citations (07/2022)

Astrid S. Martens, Johannes G. Bode, Peter C. Heinrich and Lutz Graeve*

Institute of Biochemistry, Universitätsklinikum der Rheinisch-Westfälischen Technischen Hochschule, Pauwelsstrasse 30, 52074 Aachen, Germany

*Author for correspondence (e-mail: lutz.graeve@post.rwth-aachen.de)


Accepted 1 August; published on WWW 4 October 2000



Astrid Martens





Johannes Bode




Lutz Graeve

Polar expression of gp80wt and deletion mutants in MDCK cells




Basolateral expression of gp80wt in stably transfected MDCK cells


xy



basolateral marker




gp80wt




overlay

xz




Apical expression of gp80 Δ CD in stably transfected MDCK cells


xy



basolateral marker




gp80 Δ CD



overlay

xz




Outline: Interleukin-6 signal transduction and its regulation

Part 1: Molecular mechanisms of IL-6 signal transduction

Part 2: Regulation of IL-6 signal transduction

- Half-lives of the signaling components
- Polar expression of IL-6 receptor- α (gp80)
- **IL-6 receptor- α shedding**
- Internalization of the ligand/IL-6 receptor complex (desensitisation)
- Phosphotyrosine-phosphatase SHP2 deactivates receptors, JAKs and STAT3
- SOCS proteins (suppressors of cytokine signaling) are feedback inhibitors
- Protein inhibitors of activated STATs (PIAS)
- Cross-talks between JAK/STAT signaling and proinflammatory cytokines (IL-1, TNF α)
 - STAT3 and NF- κ B compete for an overlapping response element
 - IL-1 stabilizes SOCS3-mRNA
 - IL-1 accelerates the internalization of the signal transducer gp130

Regulation of IL-6 receptor-mediated signaling by limited proteolysis (shedding)




A membrane-bound protease:

ADAM17 (A Disintegrin and Metalloproteinase) $\equiv$ TACE (TNF α converting enzyme) has been identified

Regulation of IL-6 receptor-mediated signaling by limited proteolysis (shedding)


- PMA (phorbol ester)/PKC (protein kinase C)
- apoptosis
- FMLP (N-formyl-met-leu-phe)
- CRP (C-reactive protein)



A membrane-bound protease:

ADAM17 (A Disintegrin and Metalloproteinase) $\equiv$ TACE (TNF α converting enzyme) has been identified

Agonistic action of the complex of IL-6/soluble-IL-6-receptor and trans-signaling between IL-6-responsive and non-responsive cells



Rose-John & Heinrich (1994), *Biochem. J.*, 300, 281-90

[671 citations](#) (07/2022)

The shedding protease is activated by PKC

Müllberg et al. (1993), *Eur. J. Immunol.*, 23, 473-80

[433 citations](#) (07/2022)

Müllberg et al. (1994), *J. Immunol.*, 152, 4958-68

[237 citations](#) (07/2022)

cleavage site

Increased IL-6 and soluble IL-6 receptor- α levels are found in serum and synovial fluid from patients with rheumatoid arthritis (RA)

Serum				
	IL-6 (ng/ml)	sol IL-6R α (ng/ml)	sol gp130 (ng/ml)	refs
Control	28	28	345	Keul et al. (1998)
RA	78	38	375	Keul et al. (1998)
	51	29	251	Richards et al. (2006)
Synovial fluid				
RA	6.4	15	250	Richards et al. (2006)

1998

Soluble IL-6 Receptor Potentiates the Antagonistic Activity of Soluble gp130 on IL-6 Responses¹

130 citations (07/2022)

Gerhard Müller-Newen,* Andrea Küster,* Ulrike Hemmann,* Radovan Keul,* Ursula Horsten,* Astrid Martens,* Lutz Graeve,* John Wijdenes,[†] and Peter C. Heinrich^{2*}

Soluble receptors for several cytokines have been detected in body fluids and are believed to modulate the cytokine response by binding the ligand and thereby reducing its bioavailability. In the case of IL-6, the situation is more complex. The receptor consists of two components, including a ligand-binding α -subunit (IL-6R, gp80, or CD126), which in its soluble (s) form (sIL-6R) acts agonistically by making the ligand accessible to the second subunit, the signal transducer gp130 (CD130). Soluble forms of both receptor subunits are present in human blood. Gel filtration of iodinated IL-6 that had been incubated with human serum revealed that IL-6 is partially trapped in IL-6/sIL-6R/sgp130 ternary complexes. sgp130 from human plasma was enriched by immunoaffinity chromatography and identified as a 100-kDa protein. Functionally equivalent rsgp130 was produced in baculovirus-infected insect cells to study its antagonistic potential on four different cell types. It was found that in situations in which cells lacking membrane-bound IL-6R were stimulated with IL-6/sIL-6R complexes, sgp130 was a much more potent antagonist than it was on IL-6R-positive cells stimulated with IL-6 alone. In the latter case, the neutralizing activity of sgp130 could be markedly enhanced by addition of sIL-6R. As a consequence of these findings, sIL-6R of human plasma must be regarded as an antagonistic molecule that enhances the inhibitory activity of sgp130. Furthermore, in combination with sIL-6R, sgp130 is a promising candidate for the development of IL-6 antagonists. *The Journal of Immunology*, 1998, 161: 6347–6355.

Outline: Interleukin-6 signal transduction and its regulation

Part 1: Molecular mechanisms of IL-6 signal transduction

Part 2: Regulation of IL-6 signal transduction

- Half-lives of the signaling components
- Polar expression of IL-6 receptor- α (gp80)
- IL-6 receptor- α shedding
- **Internalization of the ligand/IL-6 receptor complex (desensitization)**
- Phosphotyrosine-phosphatase SHP2 deactivates receptors, JAKs and STAT3
- SOCS proteins (suppressors of cytokine signaling) are feedback inhibitors
- Protein inhibitors of activated STATs (PIAS)
- Cross-talks between JAK/STAT signaling and proinflammatory cytokines (IL-1, TNF α)
 - STAT3 and NF- κ B compete for an overlapping response element
 - IL-1 stabilizes SOCS3-mRNA
 - IL-1 accelerates the internalization of the signal transducer gp130

A Di-leucine Motif and an Upstream Serine in the Interleukin-6 (IL-6) Signal Transducer gp130 Mediate Ligand-induced Endocytosis and Down-regulation of the IL-6 Receptor*

145 citations (07/2022)


(Received for publication, August 31, 1995, and in revised form, December 4, 1995)

Elke Dittrich‡, Carol Renfrew Haft§, Leon Muys¶, Peter C. Heinrich‡, and Lutz Graeve‡||

From the ‡Institute of Biochemistry, Rheinisch-Westfälische Technische Hochschule Aachen, 52057 Aachen, Germany, the §Diabetes Branch, NIDDKD, National Institutes of Health, Bethesda, Maryland 20892-1770, and the ¶Institute of Pathology, Rheinisch-Westfälische Technische Hochschule Aachen, 52057 Aachen, Germany




Elke Dittrich




Lutz Graeve

Intracellular domain of gp130



tm transmembrane domain

Internalization of monoclonal antibodies B-R3 and B-S12 by
COS-7 cells expressing gp130AA or gp130




Outline: Interleukin-6 signal transduction and its regulation

Part 1: Molecular mechanisms of IL-6 signal transduction


Part 2: Regulation of IL-6 signal transduction

- Half-lives of the signaling components
- Polar expression of IL-6 receptor- α (gp80)
- IL-6 receptor- α shedding
- Internalization of the ligand/IL-6 receptor complex (desensitization)
- **Phosphotyrosine-phosphatase SHP2 deactivates receptors, JAKs and STAT3**
- SOCS proteins (suppressors of cytokine signaling) are feedback inhibitors
- Protein inhibitors of activated STATs (PIAS)
- Cross-talks between JAK/STAT signaling and proinflammatory cytokines (IL-1, TNF α)
 - STAT3 and NF- κ B compete for an overlapping response element
 - IL-1 stabilizes SOCS3-mRNA
 - IL-1 accelerates the internalization of the signal transducer gp130


The IL-6 / JAK / STAT signaling cascade is well controlled



The IL-6 / JAK / STAT signaling cascade is well controlled




The IL-6 / JAK / STAT signaling cascade is well controlled



Inhibition of IL-6 signaling through SHP2

SHP2 = SH2-containing Phosphatase 2



Schaper et al. *Biochem J* 1998

141 citations (07/2022)

Anhuf et al. *J Immunol* 2000

41 citations (07/2022)

Lehmann et al. *J Biol Chem* 2003


184 citations (07/2022)

Clahsen et al. *Cell Signal* 2005

16 citations (07/2022)

Inhibition of IL-6 signaling through SHP2

SHP2 = SH2-containing Phosphatase 2



Schaper et al. *Biochem J* 1998

141 citations (07/2022)

Anhuf et al. *J Immunol* 2000

41 citations (07/2022)

Lehmann et al. *J Biol Chem* 2003


184 citations (07/2022)

Clahsen et al. *Cell Signal* 2005


16 citations (07/2022)

Inhibition of IL-6 signaling through SHP2

SHP2 = *SH2-containing Phosphatase 2*




Receptor-targeted SHP2 counteracts gp130-dependent gene induction




HepG2


SHP2 counteracts gp130-mediated signal transduction



murine embryonal fibroblasts



Affinities of SOCS3 and SHP2 to murine pY757-gp130 peptide



	K_D [nM]
SHP2-SH2 (N)	1200
SHP2-SH2 (C)	550
SHP2-SH2 (N) + SH2 (C)	170
SOCS3 full length	42
SOCS3-SH2-domain	140

KIR Kinase Inhibitory Region
SH2 Src Homology 2
ESS Extended SH2 Subdomain
N N-terminal
C C-terminal

Outline: Interleukin-6 signal transduction and its regulation

Part 1: Molecular mechanisms of IL-6 signal transduction

Part 2: Regulation of IL-6 signal transduction

- Half-lives of the signaling components
- Polar expression of IL-6 receptor- α (gp80)
- IL-6 receptor- α shedding
- Internalization of the ligand/IL-6 receptor complex (desensitization)
- Phosphotyrosine-phosphatase SHP2 deactivates receptors, JAKs and STAT3
- **SOCS proteins (suppressors of cytokine signaling) are feedback inhibitors**
- Protein inhibitors of activated STATs (PIAS)
- Cross-talks between JAK/STAT signaling and proinflammatory cytokines (IL-1, TNF α)
 - STAT3 and NF- κ B compete for an overlapping response element
 - IL-1 stabilizes SOCS3-mRNA
 - IL-1 accelerates the internalization of the signal transducer gp130

1997

SOCS proteins (suppressors of cytokine signaling) were independently discovered in 3 different laboratories

A new protein containing an SH2 domain that inhibits JAK kinases

Takaho A. Endo^{*,†}, Masaaki Masuhara^{*,†}, Masahiro Yokouchi^{*,†,‡}, Ritsu Suzuki^{*,‡}, Hiroshi Sakamoto^{*}, Kaoru Mitsui^{*}, Akira Matsumoto^{*}, Shyu Tanimura^{*}, Motoaki Ohtsubo^{*}, Hiroyuki Misawa^{*}, Tadaaki Miyazaki[§], Nogueira Leonor[§], Tadatsugu Taniguchi[§], Takashi Fujita^{||}, Yuzuru Kanakura[†], Seturo Komiya[‡] & Akihiko Yoshimura^{*}

^{*} Institute of Life Science, and [‡] Department of Orthopedic Surgery, Kurume University, Aikawamachi 2432-3 Kurume 839, Japan

[§] Department of Immunology, Faculty of Medicine, University of Tokyo, Bunkyo-ku, Tokyo 113, Japan

Nature volume 387, pages 921–924
Citations 1184 (07/2022)

A family of cytokine-inducible inhibitors of signalling

Robyn Starr^{*}, Tracy A. Willson^{*}, Elizabeth M. Viney^{*}, Leecia J. L. Murray^{*}, John R. Rayner[†], Brendan J. Jenkins[†], Thomas J. Gonda[†], Warren S. Alexander^{*}, Donald Metcalf^{*}, Nicos A. Nicola^{*} & Douglas J. Hilton^{*}

^{*} The Walter and Eliza Hall Institute for Medical Research and The Cooperative Research Center for Cellular Growth Factors, Parkville, Victoria, Australia 3052

[†] The Hanson Centre for Cancer Research, IMVS, Adelaide, Southern Australia, Australia 5000

Nature volume 387, pages 917–921
Citations 1727 (07/2022)


Structure and function of a new STAT-induced STAT inhibitor

Tetsuji Naka^{*}, Masashi Narazaki^{*}, Moritoshi Hirata^{*}, Tomoshige Matsumoto^{*}, Seiji Minamoto^{*}, Atsufumi Aono^{*}, Norihiro Nishimoto[†], Tadahiro Kajita[‡], Tetsuya Taga[§], Kazuyuki Yoshizaki[†], Shizuo Akira^{||} & Tadamitsu Kishimoto^{*}

^{*} Osaka University Medical School Department of Medicine III. 2-2, Yamada-Oka, Suita, Osaka 565, Japan

Nature volume 387, pages 924–929
Citations 1096 (07/2022)

The SOCS (SUPPRESSORS OF CYTOKINE SIGNALING) family




SOCS=Suppressor of Cytokine Signaling



Schmitz et al. *J Biol Chem* 2000
Friederichs et al. *Eur J Biochem* 2001
Fischer et. al. *Biochem J* 2004

Inhibition through SOCS-*feedback* inhibitors

SOCS=Suppressor of Cytokine Signaling



Schmitz et al. *J Biol Chem* 2000
Friederichs et al. *Eur J Biochem* 2001
Fischer et. al. *Biochem J* 2004


319 citations (07/2022)

23 citations (07/2022)

58 citations (07/2022)


Inhibition through SOCS-*feedback* inhibitors

SOCS=Suppressor of Cytokine Signaling





- | | | |
|--|---------------|-----------|
| Schmitz et al. <i>J Biol Chem</i> 2000 | 319 citations | (07/2022) |
| Friederichs et al. <i>Eur J Biochem</i> 2001 | 23 citations | (07/2022) |
| Fischer et. al. <i>Biochem J</i> 2004 | 58 citations | (07/2022) |

SOCS=Suppressor of Cytokine Signalling



SOCS3 binds the phosphotyrosine motif 759 of gp130

binding to gp130 phospho-peptides (13-mers)




COS7 cells

- SOCS1 inhibits *via* direct binding to JAKs
- SOCS3 inhibits *via* receptor binding


SOCS3-mediated signal attenuation is SHP2-independent

gene induction

SHP2 wt



SHP2 mut



SOCS3 Exerts Its Inhibitory Function on Interleukin-6 Signal Transduction through the SHP2 Recruitment Site of gp130*

(Received for publication, May 27, 1999, and in revised form, January 19, 2000)

Jochen Schmitz, Manuela Weissenbach, Serge Haan, Peter C. Heinrich†, and Fred Schaper

From the Institut für Biochemie, Rheinisch-Westfälische Technische Hochschule Aachen, Pauwelsstraße 30, D-52074 Aachen, Germany

319 citations (07/2022)

SHP2 and SOCS3 Contribute to Tyr-759-dependent Attenuation of Interleukin-6 Signaling through gp130*

Received for publication, October 15, 2002

Published, JBC Papers in Press, October 27, 2002, DOI 10.1074/jbc.M210552200

Ute Lehmann‡§, Jochen Schmitz‡§, Manuela Weissenbach‡, Radoslaw M. Sobota‡, Michael Hörtnert¶, Kerstin Friederichs‡, Iris Behrmann‡, William Tsiaris||, Atsuo Sasaki, Jens Schneider-Mergener‡‡, Akihiko Yoshimura**, Benjamin G. Neel||, Peter C. Heinrich‡§§, and Fred Schaper‡§§**

Jochen Schmitz as a post-doc at DNAX cloned IL-33

184 citations (07/2022)



Jochen Schmitz



Ute Lehmann



Manuela Decker



Serge Haan




Radek Sobota




Fred Schaper


The JAK/STAT signalling cascade is well controlled



The JAK/STAT signalling cascade is well controlled



The JAK/STAT signalling cascade is well controlled




Outline: Interleukin-6 signal transduction and its regulation

Part 1: Molecular mechanisms of IL-6 signal transduction

Part 2: Regulation of IL-6 signal transduction

- Half-lives of the signaling components
- Polar expression of IL-6 receptor- α (gp80)
- IL-6 receptor- α shedding
- Internalization of the ligand/IL-6 receptor complex (desensitization)
- Phosphotyrosine-phosphatase SHP2 deactivates receptors, JAKs and STAT3
- SOCS proteins (suppressors of cytokine signaling) are feedback inhibitors
- **Protein inhibitors of activated STATs (PIAS)**
- Cross-talks between JAK/STAT signaling and proinflammatory cytokines (IL-1, TNF α)
 - STAT3 and NF- κ B compete for an overlapping response element
 - IL-1 stabilizes SOCS3-mRNA
 - IL-1 accelerates the internalization of the signal transducer gp130

PIAS3 exerts a profound inhibitory effect on STAT3-mediated transcription of target promoters. The zinc finger protein Gfi-1 can overcome the PIAS3 block.



The EMBO Journal Vol. 19 No. 21 pp. 5845–5855, 2000

2000

The zinc finger protein Gfi-1 can enhance STAT3 signaling by interacting with the STAT3 inhibitor PIAS3

114 citations (07/2022)



**Bernd Rödel, Kamiab Tavassoli,
Holger Karsunky, Thorsten Schmidt¹,
Malte Bachmann, Fred Schaper²,
Peter Heinrich², Ke Shuai³,
Hans Peter Elsässer⁴ and Tarik Möröy⁵**

Institut für Zellbiologie (Tumorforschung), IFZ, Universitätsklinikum
Essen, Virchowstrasse 173, D-45122 Essen, ¹Bayer AG,
Apratherweg 18a, D-42096 Wuppertal, ²Institut für Biochemie,
Universitätsklinikum, RWTH Aachen, Pauwelsstrasse 30,
D-52074 Aachen, ⁴Institut für Zytobiologie und Zytopathologie,
Philipps Universität Marburg, D-35033 Marburg, Germany and
³Division of Hematology/Oncology, UCLA School of Medicine,
Los Angeles, CA, USA

⁵Corresponding author
e-mail: moeroey@uni-essen.de

B.Rödel and K.Tavassoli contributed equally to this work

The Zn-finger protein Gfi relieves the PIAS3-mediated block of STAT3-induced α 1-antichymotrypsin gene activation




Outline: Interleukin-6 signal transduction and its regulation

Part 1: Molecular mechanisms of IL-6 signal transduction



Part 2: Regulation of IL-6 signal transduction

- Half-lives of the signaling components
- Polar expression of IL-6 receptor- α (gp80)
- IL-6 receptor- α shedding
- Internalization of the ligand/IL-6 receptor complex (desensitization)
- Phosphotyrosine-phosphatase SHP2 deactivates receptors, JAKs and STAT3
- SOCS proteins (suppressors of cytokine signaling) are feedback inhibitors
- Protein inhibitors of activated STATs (PIAS)
- **Cross-talks between JAK/STAT signaling and proinflammatory cytokines (IL-1, TNF α)**
 - **STAT3 and NF- κ B compete for an overlapping response element**
 - **IL-1 stabilizes SOCS3-mRNA**
 - **IL-1 accelerates the internalization of the signal transducer gp130**


IL-1 β inhibits IL-6-induced α_2 macroglobulin synthesis in rat hepatocytes




How does IL-1 β affect IL-6-mediated signaling?





How does IL-1 β affect IL-6-mediated signaling?




Interleukin-1-signal transduction



IL-1 β inhibits α_2 M-promoter-activity via NF- κ B activation




HepG2






NF- κ B and STAT3 bind to the α_2 M-promoter

anti-p65	-	-	-	+	-	-	+
anti-STAT3	-	-	+	-	-	+	-
IL-6	-	+	+	+	+	+	+
IL-1 β	-	-	-	-	+	+	+




IL-6	-	+	+	+	+	+	+
p50	-	-	[Red triangle indicating increasing concentration]				




NF- κ B and STAT3 bind to the α_2 M-promoter



anti-p65	-	-	-	+	-	-	+
anti-STAT3	-	-	+	-	-	+	-
IL-6	-	+	+	+	+	+	+
IL-1 β	-	-	-	-	+	+	+




IL-6	-	+	+	+	+	+	+
p50	-	-					



HepG2



The α_2 -Macroglobulin Promotor




STAT3 and NF- κ B compete for an overlapping response element

The Inhibitory Effect of IL-1 β on IL-6-Induced α_2 -Macroglobulin Expression Is Due to Activation of NF- κ B¹


Journal of Immunology, 2001, 167:1469-1481

Johannes G. Bode,^{2*} Richard Fischer,[†] Dieter Häussinger,[†] Lutz Graeve,^{*} Peter C. Heinrich,^{3*} and Fred Schaper^{*} [43 citations](#) (07/2022)


Summary so far: regulation of IL-6-signaling by IL-1 β




Summary so far: regulation of IL-6-signaling by IL-1 β




Summary so far: regulation of IL-6-signaling by IL-1 β



Summary so far: regulation of IL-6-signaling by IL-1 β




Summary so far: regulation of IL-6-signaling by IL-1 β



(2) IL-1 enhances SOCS3-transcription through an unknown protein

Summary so far: regulation of IL-6-signaling by IL-1 β



(3) IL-1 stabilizes SOCS3-mRNA and enhances STAT3 inhibition


Regulation of IL-6 signaling by IL-1/TNF α : Involvement of the MKK6/p38/MK2 cascade and TTP-mediated SOCS3-mRNA stability

Ehlting et al. (2007) J. Immunol. 178, 2813-2826

88 citations (07/2022)

Bode et al. (2001) Biol Chem 382, 1447- 1453

53 citations (07/2022)




Regulation of IL-6 signaling by IL-1/TNF α : Involvement of the MKK6/p38/MK2 cascade and TTP-mediated SOCS3-mRNA stability

Ehlting et al. (2007) J. Immunol. 178, 2813-2826

88 citations (07/2022)

Bode et al. (2001) Biol Chem 382, 1447- 1453

53 citations (07/2022)




Regulation of IL-6 signaling by IL-1/TNF α : Involvement of the MKK6/p38/MK2 cascade and TTP-mediated SOCS3-mRNA stability

Ehlting et al. (2007) J. Immunol. 178, 2813-2826

88 citations (07/2022)

Bode et al. (2001) Biol Chem 382, 1447- 1453

53 citations (07/2022)




Regulation of IL-6 signaling by IL-1/TNF α : Involvement of the MKK6/p38/MK2 cascade and TTP-mediated SOCS3-mRNA stability

Ehlting et al. (2007) J. Immunol. 178, 2813-2826


88 citations (07/2022)

Bode et al. (2001) Biol Chem 382, 1447- 1453


53 citations (07/2022)




Pro-inflammatory cytokines restrict IL-6 signaling through serine-phosphorylation of gp130, internalisation and degradation




Pro-inflammatory cytokines restrict IL-6 signaling through serine-phosphorylation of gp130, internalisation and degradation



Pro-inflammatory cytokines restrict IL-6 signaling through serine-phosphorylation of gp130, internalisation and degradation



Pro-inflammatory cytokines restrict IL-6 signaling through serine-phosphorylation of gp130, internalisation and degradation



2010

Cross-regulation of cytokine signalling: pro-inflammatory cytokines restrict IL-6 signalling through receptor internalisation and degradation

76 citations (07/2022)

Simone Radtke^{1,*}, Stefan Wüller^{1,2}, Xiang-ping Yang^{1,‡}, Barbara E. Lippok¹, Barbara Mütze¹, Christine Mais³, Hildegard Schmitz-Van de Leur¹, Johannes G. Bode⁴, Matthias Gaestel⁵, Peter C. Heinrich^{1,§}, Iris Behrmann^{1,¶}, Fred Schaper^{1,**,‡‡,§§} and Heike M. Hermanns^{1,3,‡‡,§§}

¹Department of Biochemistry and Molecular Biology, Medical School RWTH Aachen, 52074 Aachen, Germany

²Department of Paediatrics, Medical School RWTH Aachen, 52074 Aachen, Germany

³Rudolf Virchow Center, DFG Research Center for Experimental Biomedicine, 97080 Würzburg, Germany

⁴Department of Gastroenterology, Hepatology and Infectiology, Medical School Heinrich-Heine University, 40225 Düsseldorf, Germany

⁵Department of Biochemistry, Medical School Hannover, 30625 Hannover, Germany

*Present address: Cancer Research UK, London Research Institute, Protein Phosphorylation Laboratory, London, WC2 A3PX, UK

‡Present address: LCBS/MIBB/NIAMS/NIH, 10 Center Drive, MSC 1930 Bethesda, MD 20892-1930, USA

§Present address: Zentrum für Biochemie und Molekulare Zellbiologie (ZBMZ), Institut für Biochemie und Molekularbiologie, Albert-Ludwigs-Universität Freiburg, Stefan Meier Str. 17, 79104 Freiburg, Germany

¶Present address: Laboratoire de Biologie et Physiologie Integree (LBPI), University of Luxembourg, 162a avenue de la Faiencerie, 1511 Luxembourg, Luxembourg

**Present address: Department of Systems Biology, Otto-von-Guericke-University Magdeburg, Universitätsplatz 2, 38106 Magdeburg, Germany

‡‡These authors contributed equally to this work

§§Authors for correspondence (schaper@rwth-aachen.de; heike.hermanns@virchow.uni-wuerzburg.de)

Accepted 9 January 2010

Journal of Cell Science 123, 947-959

© 2010. Published by The Company of Biologists Ltd

doi:10.1242/jcs.065326

2003

REVIEW ARTICLE

Principles of interleukin (IL)-6-type cytokine signalling and its regulation¹

Peter C. HEINRICH², Iris BEHRMANN, Serge HAAN, Heike M. HERMANN, Gerhard MÜLLER-NEWEN and Fred SCHAPER

Institut für Biochemie, RWTH Aachen, Universitätsklinikum, Pauwelsstrasse 30, D-52074 Aachen, Germany

2303 citations (07/2022)

The IL (interleukin)-6-type cytokines IL-6, IL-11, LIF (leukaemia inhibitory factor), OSM (oncostatin M), ciliary neurotrophic factor, cardiotrophin-1 and cardiotrophin-like cytokine are an important family of mediators involved in the regulation of the acute-phase response to injury and infection. Besides their functions in inflammation and the immune response, these cytokines play also a crucial role in haematopoiesis, liver and neuronal regeneration, embryonal development and fertility. Dysregulation of IL-6-type cytokine signalling contributes to the onset and maintenance of several diseases, such as rheumatoid arthritis, inflammatory bowel disease, osteoporosis, multiple sclerosis and various types of cancer (e.g. multiple myeloma and prostate cancer). IL-6-type cytokines exert their action via the signal transducers gp (glycoprotein) 130, LIF receptor and OSM receptor leading to the activation of the JAK/STAT (Janus kinase/signal

transducer and activator of transcription) and MAPK (mitogen-activated protein kinase) cascades. This review focuses on recent progress in the understanding of the molecular mechanisms of IL-6-type cytokine signal transduction. Emphasis is put on the termination and modulation of the JAK/STAT signalling pathway mediated by tyrosine phosphatases, the SOCS (suppressor of cytokine signalling) feedback inhibitors and PIAS (protein inhibitor of activated STAT) proteins. Also the cross-talk between the JAK/STAT pathway with other signalling cascades is discussed.

Key words: cytokine signalling, glycoprotein 130 (gp 130) interleukin-6 (IL-6), Janus kinase (JAK), mitogen-activated protein kinase (MAPK), signal transducer and activator of transcription (STAT).

Many thanks to our collaborators*

Name	Location
Fernando Bazan	Palo Alto
Kurt Binder	Vienna
Markus Böhm	Münster
Anja Bosserhoff	Regensburg
Jean Content	Brussels
Karl Decker	Freiburg
Wolfgang Doppler	Innsbruck
Alan D. Elbein	Seattle
Nelson Fausto	Seattle
Matthias Gaestel	Hannover
Joachim Grötzinger	Kiel
Kevin Harris	Birmingham, US
Dieter Häussinger	Düsseldorf
Douglas Hilton	Melbourne
Toshio Hirano	Osaka
Yannick Jacques	Nantes
James Johnston	Belfast

Name	Location
Michael Kalai	Gent
Rob Kastelein	Palo Alto
Ian Kerr	London
Tatamitsu Kishimoto	Osaka
Detlef Klimpe	Aachen
Bernd Klosterhalfen	Aachen
Tomasz Kordula	Richmond
Fritz Kreuzaler	Aachen
Dieter Kube	Göttingen
Stephan Ludwig	Düsseldorf
Felix Montero	Marseille
John Mayer	Nottingham
Tarik Möröy	Essen
Werner Müller-Esterl	Mainz
Benjamin Neel	Boston
Stefan Pflanz	Palo Alto
Ulf Rapp	Würzburg

Name	Location
Gerhard Rogler	Regensburg
Richard Roth	Stanford
Rainer Rudolph	Tutzing
Jens Schneider-Mergener	Berlin
Gerhard Schreiber	Melbourne
Jan Tavernier	Gent
Thuy-Anh Tran-Thi	Freiburg
Jim Travis	Atlanta
Axel Ullrich	Martinsried
Ulrike von Rangow	Aachen
Claudia Wellbrock	London
Sabine Werner	Zurich
John Wijdenes	Besançon
Patricia Woo	London
George Yeoh	Perth
Akihiko Yoshimura	Fukuoka

* The location of some collaborators have changed.

Many young scientists were mentioned during this presentation.

I apologize that not all of our former collaborators could be cited and shown (photos).

Sincere thanks to all the master-, MD-, PhD- students and Post-docs for the very fruitful collaboration during 32 years in Freiburg and in Aachen.

Fortunately, the driving force for all of them was always curiosity, enthusiasm, intelligence and hard work.

We had many interesting years of joint efforts and fun in working together. Thank you again!

Our research activities were only possible with the continuous support from the German Research Foundation (Deutsche Forschungsgemeinschaft, DFG), the European Commission (EC) and some others

RESEARCH GRANTS

- Special Collaborative Research Center (Sonderforschungsbereich, SFB 206)
„Biological signal reaction chains“ (1983 – 1987) Institute of Biology II, University of Freiburg
Coordinator Eberhard Schäfer, Principal investigator: Peter C. Heinrich, Freiburg
- EC-Project „Proteinases and inhibitors“
Principal investigators: John Mayer, Nottingham(UK) and Peter C. Heinrich, Freiburg
- Core Program (DFG-Schwerpunktprogramm)
„Signaltransduction in biological membranes“ (1988 – 1994, coordinator Günter Schultz, Berlin)
- DFG-Research Unit (DFG-Forschergruppe)
„Molecular mechanisms of inflammation: cytokine signaling and transport“
(1994 – 1999, coordinators Peter C. Heinrich and Siegfried Matern, RWTH Aachen)

RESEARCH GRANTS cont.

- Special Collaborative Research Center (Sonderforschungsbereich, SFB 542)

„Molecular mechanisms of cytokine-mediated inflammatory processes:
signal transduction and pathophysiological consequences”

(1999 – 2007), coordinator Peter C. Heinrich, Dept. of Biochemistry, RWTH Aachen

Forschungsberichte 1997 & 1998, 1999 & 2000, 2001 & 2002, 2003 & 2004, 2005, 2006, 2007

- EC-Project (1996 – 2000) „IL-11/receptor-complex: rational design of agonists/antagonists and immunoassays potentially used in human therapy“

Five research groups:

- Unite INSERM Institut de Biologie Nantes (Yannick Jaques, Coordinator),
- Institut Pasteur de Bruxelles (Jean Content)
- ImmunoTech S.A. Marseille (Félix Montero),
- Institute of Biochemistry, RWTH Aachen (Joachim Grötzinger, Axel Wollmer),
- Institute of Biochemistry, RWTH Aachen (Peter C. Heinrich and Gerhard Müller-Newen)

Three years support


- Interdisciplinary center for clinical research „BIOMAT“ (Biomaterials)

„Intracellular IL-6 signal transduction: pathogenic relevance of IL-6 for β 2-microglobulin amyloidosis as a consequence of bioincompatibility of dialysis membranes“

RESEARCH GRANTS cont.

- BMFT Bundesministerium für Forschung und Technologie (Federal Ministry of Research and Technology)
Projektträgerschaft Forschung im Dienste der Gesundheit
“Human chondrocytes, synovial cells, endothelial cells and monocytes/macrophages as a location of synthesis and as target cells for IL-1, IL-6 and TNF”
Joint project of the institutes of Clinical Chemistry and Biochemistry, RWTH Aachen
I UM 8916/9 (Helmut Greiling and Peter. C. Heinrich principal investigators)
- Hermann und Lilly Schilling Stiftung: Friedemann Horn, endowed professorship (Stiftungsprofessur),
07/96 - 04/97
- Volkswagenstiftung (Volkswagen foundation)
 - Marcin Wiznerowicz, Poznan , 12/94
 - Marcin Bugno, Krakau, 07/94 – 9/94
 - Marcin Kortylewski, Poznan , 10/96 –10/98
- DAAD (Deutscher Akademischer Austauschdienst)
 - Elke Roeb

Molecular mechanisms of inflammation: cytokines, signal transduction and transport






DFG



**Special collaborative research center
(Sonderforschungsbereich 542)
1999 - 2011**

**Molecular mechanisms of cytokine-mediated inflammatory processes:
signal transduction and pathophysiological consequences**

Cytokines and Receptors



Behrmann (B1)

Schaper (B2)

Schaper/Heinrich (B7)


Hermanns/Heinrich (B6)

Bode/Schaper/
Heinrich/
Häussinger (B4)


Lückhoff (B5)

Becker/Joost (B3)

Lüscher (B8)



Signal Transduction



Pathophysiological Consequences



SFB 1999